

SIG Codes

The following SIG codes are standard in Propel Rx. Additional codes can be customized in List Maintenance (visit <u>Propel Rx Online Help</u> for instructions).

sw	Shake Well	РО	By Mouth	нѕ	At Bedtime
		AU	In Each Ear	STAT	At Once
APP	Apply	OU	In Each Eye	DY	Daily
CR	Crush	IEN	In Each Nostril	D	Day
DR	Drink	AL	In the Left Ear	DS	Days
G	Give	AD	In the Right Ear	QD	Each Day
I	Inhale	os	In the Left Eye	QPM	Each Evening
INJ	Inject	OD	In the Right Eye	QAM	Each Morning
IR	Insert	R	Into the Rectum	Q	Every
INS	Instil	V	Into the Vagina	Q1H	Every Hour
PL	Place	AA	To Affected Area	Q2D	Every Second Day
T	Take	SL	Under the Tongue	Q2H	Every 2 Hours
				Q3H	Every 3 Hours
SS	One Half	PC	After Meals	Q4H	Every 4 Hours
		AC	Before Meals	Q6H	Every 6 Hours
APL	Applicatorful	DR	Drink	Q8H	Every 8 Hours
CAP	Capsule	FL	Fluids	Q12H	Every 12 Hours
CAPS	Capsules	JU	Juice	Q2-3H	Every 2-3 Hours
GTT	Drop	AQ	Water	Q2-4H	Every 2-4 Hours
GTTS	Drops	С	With	Q4-6H	Every 4-6 Hours
PF	Puffs	CF	With Food	F	For
SUPP	Suppository	CC	With Meals	QID	Four Times Daily
TBL	Tablespoon	S	Without	Н	Hour
TBLS	Tablespoonfuls			HRS	Hours
TAB	Tablet	AR	Arthritis	N	Night
TABS	Tablets	BP	Blood Pressure	QBID	One or Two Times Daily
TSP	Teaspoon	ВМ	Bowel Movement	TQID	Three or Four Times Daily
TSPS	Teaspoonfuls	CI	Circulation	TID	Three Times Daily
		CON	Constipation	BTID	Two to Three Times Daily
UD	As Directed	DI	Diarrhea	BID	Twice Daily
SP	Sparingly	FE	Fever		
ADMX	To a Maximum of	HD	Headache	MEDCP	Consultation Provided By
UF	Until Finished	HR	Heart	MEDAP	Injection Administered By
PRN	When Required	INF	Inflammation		Pharmacist
PRNF	When Necessary For	PA	Pain	MEDRA	Refer to Adaptation Form
		RA	Rash	MEDRO	Refer to Opinions Form
		SB	Shortness of Breath		

